



# Climate Report for Dumfries and Galloway

Generated on: 16/08/2026

## Introduction

This Climate Report provides high level, non-technical summaries of climate change projections for a local authority area. It uses scientific research to provide robust climate information to help decision makers plan for the future, enabling local authorities to become more resilient to climate change.

Each local authority experiences its own unique challenges from climate change. For example, urban areas are affected by the urban heat island effect resulting in higher urban temperatures compared with rural surroundings, whereas low-lying coastal areas may be at greater risk of flooding from rising sea levels.



## What affects the region's weather?

Dumfries and Galloway is located within the Western Scotland climate region. The types of weather that Western Scotland experiences across a year include:



Western Scotland has a strong maritime influence as the prevailing winds blow from the sea, providing a relatively mild climate - the mean annual temperature is 8.3°C. Mean annual temperatures are higher in coastal areas (9 to 10°C) compared to cooler upland areas (6 to 7°C).



July and August are the warmest months in the region with mean daily maxima reaching up to 19.5 °C in southern Dumfries and Galloway and the Clyde valley. Instances of extreme high temperatures are rare.




Average annual rainfall totals range from less than 1000 mm in the upper Clyde valley and along the coasts of Ayrshire and Dumfries and Galloway to over 3500 mm over the higher parts of the west Highlands, approaching the maximum values found in the UK.



West Scotland is one of the more exposed areas of the UK, being close to the Atlantic. The strongest winds are associated with the passage of deep depressions, the frequency and strength of which are greater in the winter half of the year.

## How has the climate changed in Western Scotland?





Temperature Difference (°C)  
Data: HadUK-Grid  
Concept: Ed Hawkins

Dumfries and Galloway is located within the Western Scotland climate region, where temperatures have increased (1884-2023), with many of the hottest years occurring in the last few decades

# Climate Change in the UK

## Observed changes

How have temperature and rainfall changed across the UK? These maps show changes in annual mean temperature (left) and rainfall (right) from 1991-2020 compared to a baseline period of 1961-1990. Temperatures have risen in all areas across the UK. Whilst some areas have become drier, more areas have become wetter. Some places that have become wetter over the year as a whole have also become drier in summer.



## Impacts

Urban, rural and coastal regions across the UK are already experiencing the impacts of climate change. The negative impacts of climate change may include:



### Heat

Increased energy demand for summer cooling



### Sea Level Rise

Increased risk of coastal flooding



### Heavy Rainfall

Increased risk of river and surface water flooding




### Health

Increased risk to health from heat stress



### Drought

Risk to water supplies from drought



### Drainage

Increased disruption to urban drainage systems



### Transport

Increased disruption to transport due to heat e.g. rail buckling



### Environment

Increased risk to biodiversity (plants and animals)



### Energy

Infrastructure such as gas pipes are at high risk from flooding events

## Future headlines

The climate is already changing, and we are already seeing impacts. But how might the UK's climate change in the future? The amount of future climate change will depend on how much greenhouse gas the world emits. However, even in the most optimistic scenario we are locked in to some further climate change.



There is an increased chance of **warmer, wetter winters and hotter, drier summers**.



Although the trend is for drier summers in the future, there may be increases in the intensity of heavy summer rainfall events.



Hot summers are expected to become more common. By 2050, every other summer may be as hot as the record breaking summer of 2018.




Sea level will continue to rise in the 21st century even if greenhouse gas emissions are reduced rapidly.


## Local changes in the global context

### Global Warming Levels

Global Warming Levels (GWLs) are a simple way to represent climate change at the global scale, which then drives local changes. They allow us to explore different strands of climate hazard information consistently. Relative to the pre-industrial baseline (1850-1900), central estimates for current global warming are 1.34 - 1.41°C.



Independent guidance, adopted by the UK government for the Climate Change Risk Assessment (CCRA) and 3rd National Adaptation Plan (NAP3), is to prepare for a 2°C rise in global temperature, whilst assessing the risks for 4°C.



The Paris Agreement says that we must limit global warming to well below 2°C, whilst aiming for 1.5°C.




Limiting warming to below 1.5°C above pre-industrial levels will require bigger emission reductions than currently pledged by nations around the world. Current emission reduction pledges, made as part of nationally determined contributions, are likely to lead to warming above 2°C.

### Timing of changes



The time when a particular GWL is reached will depend on future global emissions and the sensitivity of the climate system. The two futures present Central and High estimates of global warming over the 21st century. Both are consistent with current-policy global emissions reductions. They have been selected by the Climate Change Committee (CCC) for exploration for the fourth Climate Change Risk Assessment (CCRA4). These futures show the times when particular GWLs may be reached; 4°C may be reached by the end of the century under the High GWL future, but has a low likelihood. The uncertainty in these futures increases over the course of the 21st century.



## Local climate changes

The table shows projected **changes in climate** for the Local Authority area for a number of Global Warming Levels (GWLs). In each case there is a central projection (the Median) and an uncertainty range (the Lower and Upper values are the 10th and 90th percentiles). Changes are relative to 1981-2000.

The underlying science is explained in the Scientific Detail (QR Code).






The map shows the Local Authority area.

|                                                                                    |                            | 0.6°C GWL<br>Baseline 1981-2000 | 1.0°C GWL<br>Recent Past 2001-2020 | 1.5°C GWL<br>Paris Agreement | 2°C GWL<br>Guidance: Prepare | 4°C GWL<br>Guidance: Assess risks |
|------------------------------------------------------------------------------------|----------------------------|---------------------------------|------------------------------------|------------------------------|------------------------------|-----------------------------------|
|                                                                                    | TEMPERATURE                | °C                              | °C                                 | °C change                    | °C change                    | °C change                         |
|  | Summer Maximum Temperature | 25.3<br>24.9 to 25.7            | 26.7<br>25.6 to 27.5               | +1.6<br>+0.8 to +2.5         | +2.4<br>+1.4 to +3.9         | +5.4<br>+4.2 to +6.9              |
|  | Summer Average Temperature | 13.3<br>13.2 to 13.4            | 14.1<br>13.7 to 14.6               | +1.1<br>+0.6 to +1.8         | +1.7<br>+0.7 to +2.2         | +3.5<br>+2.8 to +4.2              |
|  | Winter Average Temperature | 3.2<br>3.1 to 3.2               | 3.7<br>3.2 to 4.2                  | +0.9<br>+0.1 to +1.3         | +1.2<br>+0.6 to +1.8         | +2.7<br>+1.6 to +3.2              |
|  | Winter Minimum Temperature | -8.4<br>-8.8 to -8.1            | -7.2<br>-8.7 to -5.9               | +1.6<br>+0.6 to +3.0         | +2.8<br>+0.9 to +3.8         | +4.8<br>+3.4 to +5.8              |
|  | Annual Average Temperature | 7.9<br>7.9 to 8.0               | 8.6<br>8.2 to 9.0                  | +0.9<br>+0.6 to +1.3         | +1.4<br>+0.8 to +1.7         | +3.0<br>+2.3 to +3.4              |
|                                                                                    | PRECIPITATION              | mm/day                          | mm/day                             | % change                     | % change                     | % change                          |
|  | Summer Precipitation Rate  | 3.20<br>3.14 to 3.25            | 3.07<br>2.71 to 3.66               | -2<br>-17 to +8              | -5<br>-18 to +4              | -28<br>-41 to -13                 |
|  | Winter Precipitation Rate  | 5.08<br>4.98 to 5.25            | 5.36<br>4.97 to 5.62               | +6<br>-2 to +18              | +10<br>-2 to +18             | +22<br>+16 to +32                 |

## Local climate indicators

The table shows projected **climate indicators** for the Local Authority area for a number of Global Warming Levels. For each these are annual totals: a central projection (the Median) and an uncertainty range (the Lower and Upper values are the 10th and 90th percentiles). See also the Scientific Detail (QR Code).



|                                                                                                                                                                                                                                                                                                                                                                      | 0.6°C GWL<br>Baseline<br>1981-2000 | 1.0°C GWL<br>Recent Past<br>2001-2020 | 1.5°C GWL<br>Paris<br>Agreement | 2°C GWL<br>Guidance:<br>Prepare | 4°C GWL<br>Guidance:<br>Assess risks |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------|---------------------------------------|---------------------------------|---------------------------------|--------------------------------------|
|  <b>Summer Days*</b><br>Daily maximum temperature > 25°C<br>High daytime temperatures with health impacts for vulnerable people at risk of hospital admission or death. Transport disruption – e.g. track buckling on railways. Can also indicate periods of increased water demand. | 3<br>2 to 3                        | 5<br>4 to 7                           | 6<br>3 to 10                    | 8<br>5 to 11                    | 18<br>13 to 22                       |
|  <b>Hot Summer Days*</b><br>Daily maximum temperature > 30°C<br>Increased heat related illnesses, hospital admissions or death. Further transport disruption – e.g. track buckling on railways, road melt. Overhead power lines become less efficient.                               | 0<br>0 to 0                        | 0<br>0 to 1                           | 0<br>0 to 1                     | 0<br>0 to 1                     | 3<br>2 to 5                          |
|  <b>Extreme Summer Days*</b><br>Daily maximum temperature > 35°C<br>Increased heat related illnesses, hospital admissions or death affecting not just the vulnerable. Further transport disruption – e.g. track buckling on railways, road melt.                                    | 0<br>0 to 0                        | 0<br>0 to 0                           | 0<br>0 to 0                     | 0<br>0 to 0                     | 0<br>0 to 1                          |
|  <b>Tropical Nights</b><br>Daily minimum temperature > 20°C<br>Health impact due to high night-time temperatures with potential for heat stress. Vulnerable people at increased risk of hospital admission or death.                                                               | 0<br>0 to 0                        | 0<br>0 to 0                           | 0<br>0 to 0                     | 0<br>0 to 0                     | 0<br>0 to 0                          |
|  <b>Frost Days</b><br>Daily minimum temperature < 0°C<br>Cold weather disruption due to higher than normal chance of ice and snow.                                                                                                                                                 | 67<br>66 to 68                     | 56<br>51 to 62                        | 51<br>45 to 59                  | 44<br>35 to 54                  | 20<br>14 to 32                       |
|  <b>Icing Days</b><br>Daily maximum temperature < 0°C<br>More extreme than frost days, so more severe cold weather impacts.                                                                                                                                                        | 3<br>3 to 3                        | 2<br>1 to 3                           | 1<br>1 to 2                     | 1<br>0 to 2                     | 0<br>0 to 1                          |
|  <b>Growing Degree Days<sup>+</sup></b><br>Daily mean temperature: °C > 5.5°C<br>Energy available for plant growth over a year. This is not a measure of season length.                                                                                                            | 1,322<br>1,309 to 1,339            | 1,485<br>1,421 to 1,599               | 1,540<br>1,479 to 1,639         | 1,665<br>1,521 to 1,746         | 2,150<br>1,970 to 2,260              |
|  <b>Heating Degree Days<sup>+</sup></b><br>Daily mean temperature: °C < 15.5°C<br>Indicator of energy demand for heating.                                                                                                                                                          | 2,808<br>2,799 to 2,847            | 2,608<br>2,509 to 2,782               | 2,522<br>2,436 to 2,656         | 2,379<br>2,316 to 2,584         | 1,929<br>1,817 to 2,143              |
|  <b>Cooling Degree Days<sup>+</sup></b><br>Daily mean temperature: °C > 22°C<br>Indicator of energy demand for cooling.                                                                                                                                                            | 5<br>5 to 6                        | 10<br>7 to 14                         | 12<br>7 to 20                   | 14<br>10 to 21                  | 42<br>31 to 51                       |

\* Summer days above the stated temperature thresholds can occur at any time of year

+ Degree Days are not a number of days, but the number of degrees the daily average temperature exceeds the threshold, each day, added up over a year.

# How to use the local climate projections

We are all at different points on our climate risk and adaptation journey. These projections may be used to build awareness, contribute to a risk assessment, or inform adaptation planning or reporting.


## Climate risk

Understanding the risks that climate change poses to your community, organisation or business is the first step to taking action. The **RISKS** from a changing climate and the potential for resultant impacts, depend on three factors:

**HAZARD:** weather and climate events which may have adverse effects. Their occurrence, duration and intensity may change due to climate change.


**EXPOSURE:** the location of people, property and other resources, relative to a hazard.

**VULNERABILITY:** the likelihood of the exposed people, property or resources suffering adverse effects from the hazard. Vulnerability is reduced by the capacity of people and places to adapt or respond to the hazard.



## Selection of hazard values

The information in this report may be used to assess some of the climate and weather **HAZARDS** your local area may face, within a risk assessment procedure. It provides climate information for a range of Global Warming Levels (GWLs). The median and an uncertainty range is provided for each climate variable. A key step is to identify which you need for your risk assessment. The **SIMPLE** approach follows independent guidance, adopted by the UK government for the Climate Change Risk Assessment (CCRA) and 3rd National Adaptation Plan (NAP3). The **ARP** approach aligns with guidance for the local authority Adaptation Reporting Power pilot. In the **CUSTOM** approach you decide on the basis of your risk appetite and the relevant time horizon.




# Sea Level Rise

## Around the UK


Sea level rise (SLR) is the primary way that coastal flood risk is expected to change in the UK in the future. Over the past 30 years, the UK sea level has been rising by 3.0-5.2mm per year, compared with 1.5mm per year in the 1990s.

Past and present emissions mean that sea levels will continue to rise. The amount of sea level rise depends on the location around the UK and increases with higher emissions scenarios.



## Dumfries and Galloway

Sea levels in the Local Authority are expected to rise over the 21st century. Although this result shows the likely range, larger increases cannot be ruled out. Those with a very low risk tolerance to sea level rise should consult EA guidance on H++, which provides an estimate beyond the likely range but within physical plausibility. The science behind these sea level projections can be found in the Scientific Detail (QR code).



## Selection of hazard values

The information in this Report may be used to assess some of the sea level **HAZARDS** your local area may face. A similar procedure may be followed as for land-based climate changes (p7).

It is not possible to provide the SIMPLE or ARP approaches for sea level rise as these projections are provided for future time periods directly, rather than for global warming levels. This is because sea levels will continue to rise long after global temperatures stabilise.

### CUSTOM approach

**1. Select the climate future**  
choose Central if you have a normal risk appetite, or High if you have a low risk appetite

**2. Select the time horizon**  
2030s or 2050s or 2080s

Read the value from the figure.

## Impacts



Flooding of coastal infrastructure and services



Saltwater intrusion of aquifers and agricultural land



Flooding of coastal communities and buildings

## Take action

### Who is this for?



This Climate Report is intended to be useful to a wide range of people and organisations in the UK:


- **Local Authority officers** with service responsibilities who need information on how climate change is affecting their area
- **Councillors** who need briefings on how climate change may affect their Local Authority area
- **The public** who need to understand how climate change may affect their neighbourhood, business or organisation
- **Leaders** who need a summary of climate change to share with stakeholders and the public when raising awareness

### #GetClimateReady

Tackling climate change will require both mitigation and adaptation. **Mitigation** reduces our carbon emissions with the goal of reaching **Net Zero**.

**Adaptation** prepares us for the impacts of climate changes to which we are already committed.

You may be a concerned citizen, or have public influence or be responsible for multi-million pound budgets.



You can play a part in protecting our planet from the worst impacts of climate change and in adapting our lives to protect ourselves from the impacts that we will see. Let's get ready for tomorrow. **#GetClimateReady**

## More information

The Local Authority Climate Service community site has more information.



The Met Office welcomes feedback on this service.



The Local Authority Climate Service team invites you to contact us at: [lacs@metoffice.gov.uk](mailto:lacs@metoffice.gov.uk). This is a Beta service.

This work has been supported by Defra as part of the commitments set out within the Third National Adaptation Programme.

Version 2.0

© Crown Copyright (Met Office) 2026